

Fluid and Cytological Analysis

“No fluid, no cells, no diagnosis”

Steven J. Bailey, DVM, DABVP (Feline Specialty)

ECats Veterinary Hospital, Michigan, USA

Disclaimer:

- I am first a general practitioner, and second a diplomate of the ABVP, certified in Feline Practice. I am not a board certified clinical pathologist
 - If your clinical acumen does not support what you think you see, then you should always yield to the expertise of ACVP Diplomate.
 - My goal is to excite you, to encourage and empower you to look more, rather than getting you in over your head.
- I am a bit old school.
 - I cut my teeth in medicine when veterinarians flailingly vaccinated dogs with feline distemper and mink enteritis vaccines in our attempts to save lives.
 - In that era one needed to be a ‘jack of all trades’ as there was not a referral center on every corner.
 - I will not suggest that any of you perform any skills beyond what was taught in veterinary school (CVM ’86).
 - Any skill must be used to maintain them, “use or lose”.

What is fun about looking at fluid and cytology?

- In health, the body is composed of predictable collections of cells and fluid.
- In disease, things change. Vessels exude fluid and cells. Chemotaxis initiates an ongoing march of inflammatory cells. Neoplasia causes a proliferation of abnormal cells.
- These cells and fluids are there for your diagnostic pleasure; you simply need to ask the right questions.
- Simple, inexpensive sampling can provide your client with your ability to prognosticate, to prescribe a treatment plan, to cure and/or alleviate suffering in your chosen patient.
- Taking advantage of these sampling techniques saves your patient’s time spent suffering in referral limbo. It minimizes your client’s expense when you can give them much needed answers, sooner.
- Collectively, those of us in the trenches have the opportunity to see and recognize much that has not been reported by those in the referral settings.
 - We see 99% of the iceberg, while the ‘dips’ get to see 1%
 - We need to pay attention to what our patients show us.
 - Avoid taking presentations at face value and keep an open mind.
 - When something speaks to you, then dig deeper. Post it on VIN. Ask Dr. Google. Keep thinking about it. Get more samples.

Supplementary in-house laboratory services, as an adjunct to your outside lab?

- Adds value to your patient care
 - Takes very little time

- Takes up very little space, 7ft of counter space if you don't count the sink and the workstation which you probably already have
- You can charge for it, and it costs next to nothing¹
 - Not counting labor, income is 5x expense for in-house services.
 - Despite the reality that the author finds reasons for doing things for free (in the interests of staff or personal CE and 'fun').
- Often allows you to intervene or start provisional pharmacological therapies sooner².
- Your pathologist will know you by name
 - Your succinct history, assessments and additional information will make cases more interesting, more real.
 - Your samples will become more time consuming, but more fun The pathologist will become more engaged in cases, knowing you want more.
- Utilizes your staff, enabling them to use skills they previously learned
 - Requires engaged team approach
- Empowers your staff to know they make a difference
- Provides an ongoing source of CE for doctors and technical staff
- Fends off practice burn out.

Reasons to consider in-house cytology & fluid analysis

It is an opportunity to establish a clinical diagnosis, relearn what you once knew, reinforce what you already know, and contribute to the CE of your coworkers (doctors, technicians, and even the receptionists).

You will miss more by **not** looking, than you will by not knowing.

Technicians maintain their skills and interests by routinely examining cytology and fluid samples. It is easier, if it is routine.

You will know beforehand if your samples are acceptable, or if you need additional samples. (Be sure to label all slides.)

It guides requests for additional stains and/or cultures.

Your pathologist and microbiologist will know you by name when you send in your samples with a tentative diagnosis already established. If you send them images of your clinical samples, they may even send you photomicrographs of their examinations in return.

Alert your laboratory about risks of infectious materials so they can insure proper infectious control protocols are used.

Cytologic examination may reveal the need for tissue biopsy now, rather than later. Many neoplasms can't be confirmed as such, without seeing architecture (e.g. LSA).

Cytology may reveal the need for immediate, aggressive care (exploratory surgery in the case of septic peritoneal effusions).

In-house cytology leads to definitive care sooner, rather than later, avoiding unnecessary morbidity and client expense.

Cytology is fun, it is exciting, and your clients will appreciate your extra efforts and happily pay for your services.

Back to Basics

- The tools you need you already have.
 - Microscope-standard light microscopy 10, 40, and 100X
 - Being able to attach a camera makes it even more fun.
 - Refractometer
 - Microhematocrit centrifuge
 - Stain:
 - Romanowsky stain: Diff-Quik is a common commercially available stain
 - Methylene Blue: most commonly used for reticulocyte counts.
 - Cell Counters:
 - Total Nucleated Cell Count (TNCC):
 - Absolute numbers of cell types are important

- You must know the TNCC of the blood or fluid for the cytology differential to make sense.
 - When you send samples out, be sure to ask for a full ‘fluid analysis’, rather than just ‘cytology’
- Automated bench top analyzers (laser or impedance flow cytometers) can do this. The results you get will usually be adequate.
 - Check with your instrument manufacturer to find out if you can harm your equipment
 - Most manufacturers will not suggest you use their point of care (POC) units for anything other than whole blood
 - We need to know if there are < 500 cells, 1,500 cells, 5, 10, 50, or 100,000 cells. This does not need to be perfect, just close.
 - Clusters, clumping, and sheets of cells, and variable cell size will effect accuracy, but some POC equipment does a good job with many effusions³.
 - Lastly, some microorganisms can be confusing to these units^{4,5}.
 - Many of us grew up using manual Unopette Systems. They are no longer available.
 - Leuko-TIC Test Kit for WBC (Unopette Replacement)ⁱ
 - 0.02ml of whole blood or fluid
 - 0.38ml of stained diluent (1:20 dilution)
 - Stains the WBC for ease of counting
 - MedixCorp.com
 - Cost is about \$1.40 per test
 - Hemocytometer
 - Dilutions of hemolyzed blood or fluid samples are pipetted into the counting chambers.
 - Simple math. Each side of the counting chamber holds 0.9uL.
 - Typically, you have a standard dilution and a standard factor to multiply by.
- Selected serum and/or fluid biochemical analysis
 - I-STAT and EPOC:
 - Electrolytes, renal values, glucose, lactate, and pH
 - Bench top and hand held analyzers can assess routine serum chemistries, including:
 - Albumin, creatinine, triglycerides, glucose, and more.
 - Beta Hydroxybutyrate (Precision Ultra) for serum ketones
 - Coag Dx (SCA2000 is the old version) for PT and PTT.

Obtaining Samples:

- Venipuncture tips:
 - In addition to routine collection methods, consider using heparinized insulin syringes for obtaining smaller samples from your cats. Easier on the cat and their veins.
 - Note: the smaller diameter of the barrel means it is MUCH harder to mix the blood as it settles. For example, have you tried to ‘mix’ the blood in a spun PCV tube? You can’t. Mixing blood in an insulin syringe barrel is a bit like shaking down a thermometer. Introduce a small air bubble and make sure it goes end-to-end, several times.
 - Tip for jugular draws: you can turn the syringe around and insert the needle in the direction of the blood flow (toward the heart). Often this can salvage a ‘bad’ vein and keep you from starting another site.
 - Be aware that anesthesia will decrease the serum proteins as well as the PCV⁶⁻⁹.
- Paracentesis techniques:
 - Hypodermic needles
 - Butterfly catheters (not preferred, they slice ‘n dice)
 - Through-the-needle jugular catheters
 - Four quadrant techniques
 - Diagnostic Peritoneal Lavage
 - The syringe is not your friend
 - Red Rubber (fenestrated) catheters
 - Commercial indwelling fenestrated catheters
 - Fenestrated (multi-holed) catheters (preferred)
 - Standard over-the-needle IV catheters can be quickly modified (in seconds) for paracentesis
 - If the cavity should be continuous (the peritoneum or hemothorax) then there is no need to use ultrasound to look for ‘pockets’
 -
- FNA of solid tissues:
 - Minimal risks¹⁰
 - Excellent sensitivity and positive predictive value¹¹
 - Neoplastic, vs. inflammatory vs need a surgical biopsy?
 - If ultrasound can show you a nodule ~ 1cm, you should be able to perform FNA.
 - Some samples will do great with a traditional FNA ‘aspirate’ (e.g. MCT).
 - Other tissues do better with a gentler ‘woodpecker’ or non-aspiration technique¹²
 - Some tissue and fluid samples can be smeared (pulled) like a blood smear.
 - Other will do better squashed
 - Others (some LSA) will cause cell rupture if touch in any manner. Better to look at the edge of a sprayed preparation.
 - A note about FNA and cytology of the liver
 - Some emphasize that results don’t always correlate

- What results you do get rarely mislead.
 - If you see fatty liver cells, a suppurative response (+/- bacteria), a lymphoplasmacytic infiltrate, or neoplastic cells then this is probably 'real'.
- Swabs: multipurpose
 - Cotton tipped applicators
 - Micro brushes minimize trauma to cells
- Touch impression
 - Touch impressions from excisional biopsy specimens
 - Touch or rolled impression from core biopsy samples
 - Impressions and/or smears from scrapings
 - Impressions of all/most surgical excisional samples
 - Pick the surface you want to 'see'
 - Cut surface of GIT biopsy?
 - Mucosal surface?
 - Deep surface of suspected deep pyoderma biopsies.
 - Impression of in situ lesions
- Artifacts
 - Ultrasound gel obscuring field of view¹³
 - Ultrasound gel causing cell changes, and lysis¹⁴
 - Glove powder
 - Formalin¹²

Criteria of malignancy (basically happy & homogenous vs ugly & angry)^{1,10,15}

- Macrocytosis
- Pleomorphism
- Anisocytosis
- Anisokaryosis
- Prominent/multiple nucleoli
- Increased N:C
- Variation in the N:C ratio
- Multiple nuclei
- Abnormal mitotic figures

Routine hematology: the manual CBC (< 10 minutes, 0.06ml of blood)

- RBC's
 - Remember, the PCV is a gold standard for measuring RBC volume
 - Be confident in your equipment & staff, work with them often.
 - 10,000 RPM for five minutes (or more, for less)¹⁶
 - Small volume (0.03ml of whole blood, Stat Spin)
 - Remember the buffy coat is typically 50% WBC and 50% platelets
 - Leukocytosis, thrombocytosis, metarubricytosis can all make the buffy coat interesting
 - HCT is always a calculated value, use it cautiously
 - RBC indices from commercial labs, in cats, don't over interpret them
 - Remember the PCV varies with age
 - Plasma Protein (total protein, total solids)
 - Reticulocytes
 - New Methylene Blue Stain
 - Absolute numbers are best (just like WBCs)
 - To calculate, you need
 - The % of aggregate reticulocytes
 - The total RBCs/ul will need to be estimate if you only have a % PCV % from a spun tube.

1) Use a ratio from this patient's past:

$$X = \frac{\text{Prior RBC count} * \text{Current PCV}}{\text{Prior PCV}}$$

2) Multiply PCV x 215,000 for an estimated RBC: (20 x 215,000 = 4.3 x 10⁹ RBC/ul)

- WBC (or TNCC)
 - The basis of enumerating the differential count
 - Automated Systems are good, but you should manually evaluate the differential
 - Leuko-TIC Test Kit for WBC (see above)
 - Hemocytometer (see above).
- Differential cell count (the Diff)
 - Similar processes for fluid (blood) and tissue
 - Diff-Quik (a Romanowsky stain)
 - The 'extreme feathered edge' is overlooked and very important
 - Thrombocytopenias in cats are over-diagnosed

- Look for grossly visible platelet (and WBC) clumping
- Some platelets are activated by EDTA
- Draw fresh samples into heparinized or citrated insulin syringes
- Assess RBC, thrombocytes, and 5 nucleated cell types
- Inclusions and or free organisms:
 - Organisms:
 - Bacteria:
 - Most bacteria are Diff-Quik positive
 - Mycobacterium sp. (not reported hematologically) have a thick, waxy (hydrophobic) cell wall and they do not stain with routine stains¹⁷.
 - The sensitivity of (tissue) cytology is comparable to bacterial culture².
 - Bacteremia:
 - Sepsis and bacteremia can be cause by many organisms
 - Anaplasma sp.
 - Protozoa:
 - Cytauxzoon felis¹⁸
 - Babesia¹⁹
 - Hepatozoon²⁰
 - Trypanasoma sp.²¹
 - Toxoplasma gondii²²⁻²⁴
 - Fungi:
 - Histoplasma capsulatum (man & dog)^{25,26}
 - Candidiasis (man)²⁷
 - Not uncommon in man, are we missing it?
 - Nematodes, filarial:
 - Dirofilaria immitis
 - Brugia malayi (tropics)²⁸.
 - Metabolic & genetic disease inclusions
 - Birman & Siamese Cats^{29,30}
 - Mannosidosis³¹
 - Mucopolysaccharidosis³¹
 - Gangliosidosis³²
 - Chediak Higashi Syndrome³³

In-house cultures: bacterial

- Mostly used for urine.
 - Included as part of our routine urine analysis
- Controversial

- BSL2 facility is required for handling cultures including:
 - Bacterial, dermatophytes (fungal), and parasites
 - Urine, tissues, hair (DTM), fecal (Tritrichomonas) and other cultures
 - Positive culture plates (all plates) should be autoclaved in special bags when discarding
- In-expensive
 - Disposable calibrated loops (10- μ l)
 - Allows for quantitative fluid cultures.
 - Blood agar plates (inexpensive)
 - No flames
- Effective
 - Positive cultures are evident in 6 to 8 hours
 - More than 95% of urine cultures are negative
 - No longer 'fear' the negative urine culture
 - Can be used screening test in young cats as well
 - Recommend morphotyping with Diff-Quik stain
 - Gram staining is discouraged, unless processing multiple samples daily
 - Catalase reaction +/- is helpful when you have cocci bacteria
 - Staph sp. vs enterococcus or streptococcus
 - Osborn's chart of significance^{34,35}
- Versatile aerobic culturing
 - Urine culture
 - FNA of any tissue (liver)
 - Gallbladder-bile
 - Fluid culture
 - Blood culture (a drop of blood spread on a plate) if suspecting septicemia
 - Impressions from surgical sample (liver, pancreas)
 - Deep surface of skin punch biopsy samples

Fluid analysis (same as for a CBC)

- You need:
 - Fluid
 - Volume, color, clarity, viscosity
 - Total Protein / Fluid Specific Gravity
 - 2.5 g/dl above and below
 - Find a table of your choosing, but none are perfect
 - TNCC
 - Cytology
 - Fluid Culture (+/-)
 - Selected biochemistries (+/-)

How to make a fenestrated catheter

Steven J. Bailey, DVM, DABVP

ECats, Michigan, USA

A fenestrated catheter is a practical and indispensable diagnostic tool that facilitates establishing a diagnosis. The fashioning of a home-made, over-the-needle, multi-holed “fenestrated” catheter is a simple technique often employed in emergency and critical care practice. The cost to create the catheter is limited to the cost of the over-the-needle IV catheter, a scalpel blade, and less than 60 seconds of effort. While the catheters can be fashioned aseptically without them, a gloved technique is recommended to avoid possible contamination. These catheters can be fashioned intraoperatively for centesis of cystic lesions. Alternatively, they could be premade, repackaged, and ethylene oxide sterilized for later use. Commercially made catheters are similarly available and can be used as well (but what fun is that?).

A variety of lengths and diameters can be used. Most commonly in our practice 16 to 18 G, 1 to 2” catheters are used. These catheters are suited for paracentesis of the thorax, pericardium, peritoneum, and retroperitoneum, as well as hepatic, renal, perirenal cysts. Paracentesis techniques may be diagnostic or therapeutic.

In addition to the terminal opening of the catheter lumen, four side holes increase the likelihood of successful fluid collection³⁶. A #15 or #12 scalpel is used to cut the wall of the catheter against the side of the stylet. The holes in the catheter should be positioned at 0, 90, 180, and 270° and each approximately 1-3mm long, 1-1.5mm wide, and spaced end-to-end at least 3mm apart. No holes should be cut within 1 cm of the hub as these will be in (or outside) the body wall. If the holes are too large or spaced too close together they may weaken the shaft of the catheter and lead to buckling during insertion or breaking during extraction. Similar to whittling a stick, the side holes are cut with a smooth, flowing movement, and a flick of the wrist to finish.

When cutting the holes into the catheter this author’s technique is to aim the catheter toward myself and cut “sharp edge to sharp point.” The intent is to have no burrs remain, but if a small burr is created it should be on the leading edge of the side-hole in the catheter. This facilitates a smooth entrance of the catheter into the body cavity. Any burrs created should ‘catch’ during a slow, controlled withdrawal of the catheter from the body wall. If resistance, the ‘catch’, is felt, the catheter should be rotated along its long axis to facilitate extraction and reduce the likelihood of breaking off a portion of the catheter in the body cavity.

Hair should be clipped and the site surgically prepped as you would for IV catheter placement. Depending on the patient, local anesthesia, sedation, and/or general anesthesia are appropriate considerations. Once the catheter and stylet are inserted into the body cavity, the stylet should remain fixed while the catheter is advanced to its hub. Resistance to advancement indicated that the catheter is not into the body cavity, or there is a burr that is catching on the tissues.

While the catheter is in the cavity it may be possible to attach a closed system for removal of large volumes of fluids. Fluid will flow by capillary action and collect in the hub of the catheter. When only small amounts of fluid are present, the fluid can be aseptically drawn from the hub into an appropriately sized syringe with needle attached. Do not attach the hub of a syringe to the catheter and create negative pressure as this will cause omentum, tissues or debris to lodge in the side holes and interfere or stop the flow of fluid altogether. If there is enough volume of fluid that it drips from the hub, then the fluid can be collected directly into a blood collection tube. If the volume is large you may be able to attach a syringe or extension tubing directly to the hub and aspirate in a conventional manner. If the fluid is a hemorrhagic fluid, the syringe or collection tube may be heparinized to prevent ongoing clot formation.

PARACENTESIS

Thoracocentesis

- Generally, the patient is kept in sternal recumbency during thoracocentesis
- Don't count ribs unless placing an indwelling thoracostomy tube
- There is no need to use ultrasound to look for pockets as each hemithoracic cavity should be a contiguous space (potential space).
- You may elevate parts of the chest to get the fluid to pool dependently during thoracocentesis.
- Aseptically prep a very small area.
- Use a local block or general anesthesia.
- An IV extension set is used so the syringe is not held against the chest.
- The three-way stop-cock is added time and expense
- If there is an effusion, rarely is there enough negative intrathoracic pressure to actually suck air in, particularly if there is an extension set attached.
- Grossly, when you collect a pleural effusion you should have a visual suspicion if the collected fluid is a: transudate, chylous, exudate, pyogranulomatous, neoplastic or hemorrhagic. Based on your suspicion ask these questions:
 - What volume should be removed?
 - Should this cat have an echocardiogram to evaluate the LA and myocardium?
 - Will furosemide help, or hurt this patient.
 - What biochemical analyses are warranted?
 - Fluid triglycerides vs serum?
 - Fluid glucose or lactate?
 - What is expected cytologically, and should the smear to be looked at concurrent with abdominocentesis (takes two minutes)?
 - Are cultures (aerobic and anaerobic) warranted based on cytology?

- Can provisional pharmacological decisions be based on the morphology of visible organisms?
- Should the client be given a poorer prognosis based on suspicion of neoplastic cells?
- Is post-centesis radiography warranted?
- If the fluid is hemorrhagic, should it be collected aseptically for possible autotransfusion?

Abdominocentesis

- Generally, the author performs abdominocentesis with the cat in lateral recumbency.
 - Local block, or general anesthesia
 - Small aseptic clip and prep 3 cm cranial to the umbilicus
 - Especially when there is a small volume of effusion do not use a syringe.
 - “Syringes suck” for abdominocentesis.
 - Use a heparinized syringe and needle to draw fluid from the catheter hub.
- Grossly, when you collect an abdominal effusion you should have a visual suspicion if the collected fluid is a: transudate, suppurative exudate, pyogranulomatous, chylous, neoplastic, bilious, or hemorrhagic effusion. When should you suspect uroperitoneum?
- What volume should be removed?
- When should you consider cardiac disease and echo?
 - Will furosemide help, or hurt this patient?
- What biochemical analyses are warranted?
 - Fluid triglycerides vs serum?
 - Fluid creatinine vs serum?
 - Fluid glucose or lactate?
 - Fluid bilirubin vs serum bilirubin?
 - Albumin/globulin ratio?
- What is expected cytologically, and should the smear to be looked concurrent with the centesis (takes two minutes)?
 - Are cultures (aerobic and anaerobic) warranted based on cytology?
 - Bacteria in an abdominal effusion suggests treatment is prompt surgery rather than medical management.
 - Can you make provisional pharmacological choices based on morphology of visible organisms?
 - For hemorrhagic effusions, are you seeing platelets that suggest iatrogenic or ongoing hemorrhage?
- Should the clients be informed of a poorer prognosis based on strong suspicion of neoplastic cells?

- Is post-centesis radiography indicated?

Retroperitoneocentesis

- Can be based on US or intraoperatively
- Chylous
- Uroretroperitoneum
- Suppurative
- Others

Pericardiocentesis³⁷⁻⁴²

- Often associated with systemic disease and other effusions
- Diagnosis is usually based on radiographic signs of an enlarged rounded heart
- Ultrasound (concurrent) examination is highly recommended
 - Cytology, aerobic and anaerobic culture
 - Volume
 - 1.5 to 95ml, not correlated to clinical signs
 - Classification:
 - Transudate (hypoalbuminemia, volume overload).
 - Modified Transudate: CHF (30 to 75%?), PPDH
 - Biliary (PPDH): two reported cases
 - Suppurative: Septic (10%)
 - Eosinophilic: ECats case
 - Pyogranulomatous: FIP (>15%)
 - Neoplastic: LSA & others (>15%)
 - Hemorrhagic: coagulopathy (10%), neoplasia, L atrial tear

Cholecystocentesis^{43,44}

- FNA of the gallbladder
- With US this is quick, easy
- Bile peritonitis and/or GB rupture is a rare sequela
- Make it routine for GIT cases

Airway Sampling

- Nasal swabs and exudates
 - Don't discard that nose wipe!
 - Blind, otoscopic, or endoscopic
 - They don't reveal all, but they don't lie
 - Inflammatory cells
 - Neoplastic cells
 - Intracellular bacteria

- Fungal disease
 - Pathogenic fungi
 - Rafts of saprophytic fungi
 - *Eucoleus boehmi* (dogs)⁴⁵
 - *Pneumonyssus caninum* (nasal mite)⁴⁶
- Nasopharyngeal flush⁴⁷
 - Washes exudates and dislodges tissues
 - Impression smears and histopathology
 - Note about a special bacterium:
 - *Simonsiella* sp. belongs in the oral cavity.
 - It is distinctive and suggests oral contamination, or a very unusual infection.
- Bronchoalveolar Lavage⁴⁸
 - Blind or endoscopically guided
 - Examine sample at the time, during the collection⁴⁹
 - Homogenize, wet prep, stained smear
 - Assess cellularity
 - Assess cilia motility
- Pulmonary FNA⁵⁰
 - Generalized and nodular disease
 - Blind, or planned based on radiographs and rib spaces
 - Ultrasound may help to localize peripheral masses

Arthrocentesis (joint aspirates)⁵¹

- Septic effusions
 - *Mycoplasma* is underappreciated, difficult to see and culture
 - In-house culturing helps
- Synovial Cysts⁵²⁻⁵⁴

1. Hodges J. Using cytology to increase small animal practice revenue. *Veterinary Clinics of North America: Small Animal Practice* 2013;43:1385-1408.
2. Clinkenbeard KD, Cowell RL, Morton RJ, et al. Diagnostic cytology: bacterial infections. *The Compendium on continuing education for the practicing veterinarian (USA)* 1995.
3. Zandecki M, Genevieve F, Gerard J, et al. Spurious counts and spurious results on haematology analysers: a review. Part II: white blood cells, red blood cells, haemoglobin, red cell indices and reticulocytes. *International journal of laboratory hematology* 2007;29:21-41.
4. Gloster ES, Strauss RA, Jimenez JF, et al. Spurious elevated platelet counts associated with bacteremia. *American journal of hematology* 1985;18:329-332.
5. Branda JA, Kratz A. Effects of yeast on automated cell counting. *American journal of clinical pathology* 2006;126:248-254.
6. Marini R, Jackson L, Esteves M, et al. Effect of isoflurane on hematologic variables in ferrets. *American Journal of Veterinary Research* 1994;55:1479-1483.
7. Dhumeaux MP, Snead EC, Epp TY, et al. Effects of a standardized anesthetic protocol on hematologic variables in healthy cats. *Journal of feline medicine and surgery* 2012;14:701-705.
8. Volpato J, Mattoso CRS, Beier SL, et al. Sedative, hematologic and hemostatic effects of dexmedetomidine–butorphanol alone or in combination with ketamine in cats. *Journal of feline medicine and surgery* 2014:1098612X14549214.
9. Hikasa Y, Kawanabe H, Takase K, et al. Comparisons of sevoflurane, isoflurane, and halothane anesthesia in spontaneously breathing cats. *Vet Surg* 1996;25:234-243.
10. Barger A. Cytology of Neoplasia An Essential Component of Diagnosis. *Today's Veterinary Journal* 2012;2:6.
11. Ghisleni G, Roccabianca P, Ceruti R, et al. Correlation between fine-needle aspiration cytology and histopathology in the evaluation of cutaneous and subcutaneous masses from dogs and cats. *Veterinary Clinical Pathology* 2006;35:24-30.
12. Moore AR. Preparation of Cytology Samples: Tricks of the Trade. *Veterinary Clinics of North America: Small Animal Practice* 2016.
13. Royer MC, Davidson DD, Dimitrov RK, et al. Ultrasound gel causes fine needle aspiration artifact? A clear choice. *Acta cytologica* 2012;56:146-154.
14. Molyneux A, Coghill S. Cell lysis due to ultrasound gel in fine needle aspirates; an important new artefact in cytology. *Cytopathology* 1994;5:41-45.
15. Alleman AR, Bain P. Diagnosing neoplasia: the cytologic criteria for malignancy. *Veterinary Medicine* 2000.
16. Guest GM, Siler VE. A centrifuge method for the determination of the volume of cells in blood. *The Journal of Laboratory and Clinical Medicine* 1934;19:757-768.
17. Rivière D, Pingret J-L, Etievant M, et al. Disseminated *Mycobacterium avium* subspecies infection in a cat. *Journal of feline medicine and surgery* 2011;13:125-128.
18. Miller J, Davis CD. Increasing frequency of feline cytauxzoonosis cases diagnosed in western Kentucky from 2001 to 2011. *Veterinary parasitology* 2013;198:205-208.
19. Hartmann K, Addie D, Belák S, et al. Babesiosis in Cats ABCD guidelines on prevention and management. *Journal of feline medicine and surgery* 2013;15:643-646.
20. Baneth G, Aroch I, Tal N, et al. Hepatozoon species infection in domestic cats: a retrospective study. *Veterinary parasitology* 1998;79:123-133.

21. Da Silva AS, Costa MM, Wolkmer P, et al. Trypanosoma evansi: hematologic changes in experimentally infected cats. *Experimental parasitology* 2009;123:31-34.
22. Silveira C, Vallochi AL, da Silva UR, et al. Toxoplasma gondii in the peripheral blood of patients with acute and chronic toxoplasmosis. *British Journal of Ophthalmology* 2010;bjo. 2008.148205.
23. Arnold SJ, Kinney MC, McCormick MS, et al. Disseminated toxoplasmosis. *Archives of pathology & laboratory medicine* 1997;121:869.
24. Barrs V, Martin P, Beatty J. Antemortem diagnosis and treatment of toxoplasmosis in two cats on cyclosporin therapy. *Australian veterinary journal* 2006;84:30-35.
25. Updesh M, Sachdeva S, Das R. Peripheral smear discloses histoplasmosis. *Blood* 2010;115:3653.
26. Clinkenbeard K, Cowell R, Tyler R. Identification of Histoplasma organisms in circulating eosinophils of a dog. *Journal of the American Veterinary Medical Association* 1988;192:217.
27. Lam S, Hsia CC. Peripheral blood candidiasis. *Blood* 2012;119:4822-4822.
28. Chansiri G, Khawsak P, Phantana S, et al. The efficacy of a single-oral-dose administration of ivermectin and diethylcarbamazine on the treatment of feline Brugia malayi. 2005.
29. Hirsch V, Cunningham T. Hereditary anomaly of neutrophil granulation in Birman cats. *American Journal of Veterinary Research* 1984;45:2170-2174.
30. Thompson J. Unusual granulation of neutrophils from Siamese cats and possible link with granulation in Birman cats. *New Zealand Veterinary Journal* 2009;57:70-70.
31. Alroy J, Freden G, Goyal V, et al. Morphology of leukocytes from cats affected with α -mannosidosis and mucopolysaccharidosis VI (MPS VI). *Veterinary Pathology Online* 1989;26:294-302.
32. Hasegawa D, Yamato O, Kobayashi M, et al. Clinical and molecular analysis of GM2 gangliosidosis in two apparent littermate kittens of the Japanese domestic cat. *Journal of Feline Medicine & Surgery* 2007;9:232-237.
33. Kahraman MM, Prieur DJ. Chediak-Higashi syndrome: Prenatal diagnosis by fetal blood examination in the feline model of the disease. *American journal of medical genetics* 1989;32:325-329.
34. Osborne CA, Lulich JP. *The Bristol Veterinary Handbook of Antimicrobial Therapy* Trenton, NJ: Veterinary Learning Systems Co; 1987;74-87.
35. Polzin DJ. Management of recurrent bacterial urinary tract infections. *The Compendium on continuing education for the practicing veterinarian (USA)* 1994.
36. Crowe DT, Crane SW. Diagnostic abdominal paracentesis and lavage in the evaluation of abdominal injuries in dogs and cats: clinical and experimental investigations. *J Am Vet Med Assoc* 1976;168:700-705.
37. Rush J, Keene B, Fox P. Pericardial disease in the cat: a retrospective evaluation of 66 cases. *Journal of the American Animal Hospital Association* 1990;26:39-46.
38. Hall DJ, Shofer F, Meier CK, et al. Pericardial effusion in cats: a retrospective study of clinical findings and outcome in 146 cats. *Journal of Veterinary Internal Medicine* 2007;21:1002-1007.
39. Less RD, Bright JM, Orton EC. Intrapericardial cyst causing cardiac tamponade in a cat. *Journal of the American Animal Hospital Association* 2000;36:115-119.
40. Scruggs SM, Bright JM. Chronic cardiac tamponade in a cat caused by an intrapericardial biliary cyst. *Journal of feline medicine and surgery* 2010;12:338-340.

41. Majoy SB, Sharp CR, Dickinson AE, et al. Septic pericarditis in a cat with pyometra. *Journal of Veterinary Emergency and critical care* 2013;23:68-76.
42. Wolff CA, Fischetti AJ, Bond B, R. What Is Your Diagnosis? Bacterial Pericardial effusion. *Journal of the American Veterinary Medical Association* 2010;236:735-736.
43. Wagner KA, Hartmann FA, Trepanier LA. Bacterial culture results from liver, gallbladder, or bile in 248 dogs and cats evaluated for hepatobiliary disease: 1998–2003. *Journal of Veterinary Internal Medicine* 2007;21:417-424.
44. Brain PH, Barrs VR, Martin P, et al. Feline cholecystitis and acute neutrophilic cholangitis: clinical findings, bacterial isolates and response to treatment in six cases. *Journal of Feline Medicine & Surgery* 2006;8:91-103.
45. Piperisova I, Neel JA, Tarigo J. What is your diagnosis? Nasal discharge from a dog. *Veterinary Clinical Pathology* 2010;39:121-122.
46. Brandt RB. Pneumonyssus caninum (nasal mite) in four golden retrievers. *The Canadian Veterinary Journal* 1988;29:741.
47. Ashbaugh EA, McKiernan BC, Miller CJ, et al. Nasal hydropulsion: a novel tumor biopsy technique. *Journal of the American Animal Hospital Association* 2011;47:312-316.
48. Reinero C. Bronchoalveolar lavage fluid collection using a blind technique. *NAVC Clin Brief* 2010:58-61.
49. Nafe LA, DeClue AE, Reinero CR. Storage alters feline bronchoalveolar lavage fluid cytological analysis. *Journal of Feline Medicine & Surgery* 2011;13:94-100.
50. DeBerry JD, Norris CR, Samii VF, et al. Correlation between fine-needle aspiration cytopathology and histopathology of the lung in dogs and cats. *Journal of the American Animal Hospital Association* 2002;38:327-336.
51. Clements D. Arthrocentesis and synovial fluid analysis in dogs and cats. *In Practice* (0263841X) 2006;28.
52. White JD, Martin P, Hudson D, et al. What is your diagnosis? Synovial Cyst. *Journal of feline medicine and surgery* 2004;6:339-344.
53. Stead A, Else R, Stead M. Synovial cysts in cats. *Journal of Small Animal Practice* 1995;36:450-454.
54. Kligman K, Kim S, Winter M, et al. What is your diagnosis? Synovial cysts. *Journal of the American Veterinary Medical Association* 2009;235:945-946.

¹ Leuko-tic, Bioanalytic GmbH, Germany. Email: office@bioanalytic.de